naturalnews.com

THEY LIED: Bombshell new research confirms that Reuters, AP misled the public about how long mRNA nanoparticles persist in the body following vaccinations

JD Heyes

4-5 minutes



3 Ways Your Dog Asks For Help



Watch Out For These Signs >>>

(Natural News) Once again, the info-gatekeepers in the mainstream media – the "fact-checkers" at the *Associated Press*, *Reuters*, etc., – have served their purpose as disinformation outlets for the Deep State regime by lying about another aspect of the Wuhan coronavirus (COVID-19) vaccines.

According to Alex Berenson, a former *New York Times* reporter and author who has been following and tracking COVID-19 research and data since the pandemic began, a little-noticed report in the highly respected, peer-reviewed journal *Cell* reveals that mRNA materials from the Moderna and Pfizer vaccines remain in lymph nodes for at least 60 days, while "free-floating spike proteins circulate at high levels in the blood after vaccination."

That's not what the info-gatekeepers have been telling us.

"Vaccinated people infected with variants of Sars-Cov-2 produce antibodies biased toward the original and now extinct variant, rather than the one that has actually infected them," Berenson noted further on his SubStack account, noting that

what he says is not a conspiracy theory.

He added: "Findings from a preprint in Cell, among the world's leading scientific journals. Almost 50 researchers worldwide collaborated on the work, which Cell released online two weeks ago. The preprint has received little attention, possibly because it discusses the potential implications of its findings only obliquely. In discussing the fact that the mRNA hamper the immune response to new variants, the researchers offer extra doses as a potential solution."



While the researchers still offer up booster vaccines as a way to "compensate for relatively decreased binding" to new variants (vaccine boosters have proven dangerous and ineffective as well, by the way), Berenson noted that, based on the research, "the preprint's findings destroy comforting fictions about the mRNA shots, including that the body quickly destroys the genetic material in the jabs, as *Reuters* and other 'fact-checkers' have long insisted."

Researchers <u>writing in *Cell*</u> note that the genetic materials actually remain in the body for up to two months – 60 days.

Prolonged detection of vaccine mRNA in LN GCs, and spike antigen in LN GCs and blood following SARS-CoV-2 mRNA vaccination

The biodistribution, quantity and persistence of vaccine mRNA and spike antigen after vaccination, and viral antigens after SARS-CoV-2 infection, are incompletely understood but are likely to be major determinants of immune responses. We performed *in situ* hybridization with control and SARS-CoV-2 vaccine mRNA-specific RNAScope probes in the core needle biopsies of the insilateral axillary LNs that were collected 7-60 days after 2nd dose of mRNA-1273 or

BNT162b2 vaccination, and detected vaccine mRNA collected in the GCs of LNs on day 7, 16, and 37 post vaccination, with lower but still appreciable specific signal at day 60 (Figures 7A - 7E). Only rare foci of vaccine mRNA were seen outside of GCs. Axillary LN core needle biopsies of non-vaccinees (n = 3) and COVID-19 patient specimens were negative for vaccine probe hybridization. Immunohistochemical staining for spike antigen in mRNA vaccinated patient LNs varied between individuals, but showed abundant spike protein in GCs 16 days post-2nd dose, with spike antigen still present as late as 60 days post-2nd dose. Spike antigen localized in a reticular pattern around the GC cells, similar to staining for follicular dendritic cell processes (Figure 7B). COVID-19 patient LNs showed lower quantities of spike antigen, but a rare GC had positive staining (Figure 7F). Immunohistochemical staining for N antigen in peribronchial LN secondary and primary follicles of COVID-19 patients (Figures 7F - 7I) was positive in 5 of the 7 patients,

The mRNA vaccines cause the body to manufacture a large amount of spike proteins, and advocates for the vaccines have always argued that the proteins then remain bound to cells where they were made, Berenson noted.

with a mean percentage of nucleocapsid-positive follicles of more than 25%.

However, according to the researchers, they found that spike protein in the blood following mRNA vaccines were present and as high as those that were naturally produced after having the COVID-19 infection (again, making the vaccines for previously sickened people worthless).

Dr. Robert Malone, inventor of the mRNA technology, discussed the production of harmful amounts of spike proteins caused by the vaccines with top podcaster Joe Rogan last month.

"The gain-of-function research behind the development of the bioweapon spike protein ultimately paved the way for the

development of a similar, but different spike protein which is currently being used as mRNA instructions in the covid vaccine. The drug companies have an ethical obligation to prove that the spike protein in their vaccine is not toxic," said Malone as reported by *Natural News*.

When he was asked about the decreasing efficacy of vaccines and boosters, Malone said he agreed that the jabs are actually leading to negative efficacy, as demonstrated in studies done by Danish researchers. As doses increase, individuals are becoming more prone to being infected by the virus.

In Israel, where citizens are being pressured into getting a fourth vaccine dose, their B and T memory cells "are being trained to attack a spike protein that has long since mutated, setting the vaccinated up for antibody-dependent enhancement, autoimmune issues and cancer," *Natural News* reported.

Malone also warned that spike protein had a propensity to cause blood clotting, which has likely led to increases in strokes, heart attacks and other maladies related to the phenomenon.

This is why this vaccine should not be mandatory – at all.

Sources include:

AlexBerenson.Substack.com

NaturalNews.com